


TEXAS

Key Messages

Temperatures in Texas have risen almost 1.5°F since the beginning of the 20th century. Historically unprecedented warming is projected during this century, with associated increases in extreme heat events.

Although projected changes in annual precipitation are uncertain, increases in extreme precipitation events are projected. Higher temperatures will increase soil moisture loss during dry spells, increasing the intensity of naturally occurring droughts.

Future changes in the number of landfalling hurricanes in Texas are difficult to project. As the climate warms, hurricane rainfall rates, storm surge height due to sea level rise, and the intensity of the strongest hurricanes are all projected to increase.



The Texas climate is characterized by hot summers and mild to cool winters. Three geographical features largely influence the state’s varied climate: the Rocky Mountains block intrusions of moist Pacific air from the west and tend to channel arctic air masses southward during the winter; the relatively flat central North American continent allows easy north and south movement of air masses; and the Gulf of Mexico serves as the primary source of moisture, which is most readily available to the eastern part of the state. As a result of these factors, the state exhibits large east-west variations in precipitation and is subject to frequent and varied extreme events, including hurricanes, tornadoes, droughts, heat waves, cold waves, and extreme precipitation. Due to rapid population growth, especially in urban areas, increased demand for limited water supplies may increase Texas’s vulnerability to naturally occurring droughts.

Temperatures in Texas have risen almost 1.5°F since the beginning of the 20th century (Figure 1). While there is no overall trend in extremely hot days (Figure 2), the number of very warm nights was particularly high during the 2010s (Figure 3). The urban heat island effect increased these occurrences in city centers. The summer of 2011 was the warmest summer on record (since 1895) and broke the state record for highest average number of days with temperatures of 100°F or more. The Dallas-Fort Worth area endured 40 consecutive days with temperatures higher than 100°F, which was the second-longest streak on

Observed and Projected Temperature Change




Figure 1: Observed and projected changes (compared to the 1901–1960 average) in near-surface air temperature for Texas. Observed data are for 1900–2020. Projected changes for 2006–2100 are from global climate models for two possible futures: one in which greenhouse gas emissions continue to increase (higher emissions) and another in which greenhouse gas emissions increase at a slower rate (lower emissions). Temperatures in Texas (orange line) have risen almost 1.5°F since the beginning of the 20th century. Shading indicates the range of annual temperatures from the set of models. Observed temperatures are generally within the envelope of model simulations of the historical period (gray shading). Historically unprecedented warming is projected during this century. Less warming is expected under a lower emissions future (the coldest end-of-century projections being about 2°F warmer than the historical average; green shading) and more warming under a higher emissions future (the hottest

years in the hottest end-of-century projections being about 10°F warmer than the hottest year in the historical record; red shading). Sources: CISESS and NOAA NCEI.

record (1899–2020). The record dry conditions contributed to the higher temperatures. Daily minimum temperatures in January typically range from about 20°F in the northern Panhandle to about 50°F near the mouth of the Rio Grande River. The annual number of entire days below freezing was well above average in the 1970s and 1980s but has since been near the long-term average (Figure 4a).

Precipitation is widely variable across Texas, with normal amounts ranging from less than 10 inches in the far west to more than 60 inches in the extreme southeast. Historically significant droughts occurred in the late 1910s, the early 1950s, and the early 2010s; the driest calendar years were 1917, 1956, and 2011 (Figure 4b). The driest consecutive 5 years was the 1952–1956 interval and the wettest was the 2015–2019 period. In the 1990s and early 2000s, the number of 3-inch extreme precipitation events was above average, and after the dry period of 2005–2014, they were well above average during the 2015–2020 period (Figure 4c). The five wettest months on record have all occurred since the year 2000, led by 9.1 inches in May 2015. Hurricane Harvey (2017) was the most destructive event in Texas history, mostly due to the unprecedented rainfall, which contributed to the second wettest month on record despite affecting only part of the state. After making landfall on August 25, Harvey slowed and was nearly stationary for several days near Houston. Rainfall exceeded 30 inches in many locations, and a few locations had more than 50 inches (Figure 5). Catastrophic flooding occurred across much of southeast Texas.

Texas is consistently ranked in the top 10 states affected by extreme events. In 2020, the state was hit by eleven of the nation’s billion-dollar disasters. The three most impactful events were drought, extreme heat, and wildfires. The warmest and the driest summer in the historical record helped fuel the worst wildfire season since statewide records began (approximately 1990), with nearly 4 million acres burned and almost \$750 million in damages. Since the creation of the United States Drought Monitor Map in 2000, Texas has been completely drought-free for approximately 8% of the time (2000–2014), and at least half of the state has been under drought conditions for approximately 42% of the same period. Paleoclimatic records indicate that droughts as severe as the one in 2011 have occurred occasionally in the past 1000 years (Figure 6). Higher temperatures and drought conditions are likely to increase the severity, frequency, and extent of wildfires in the future, threatening significant harm to property, human health, and the livelihood of residents.

Over the period of 1900 to 2020, Texas endured more than 85 tropical storms and hurricanes (about 3 storms every 4 years); approximately half of them (46) were hurricanes (Figure 4d). Since 2000, Texas has experienced 19 named storms, including 8 destructive hurricanes, with Hurricane Harvey (Category 4), Hurricane Rita (Category 3), and Hurricane Ike (Category 2) causing the most significant damage. While Hurricane Rita

Observed Number of Extremely Hot Days




Figure 2: Observed annual number of extremely hot days (maximum temperature of 100°F or higher) for Texas from 1900 to 2020. Dots show annual values. Bars show averages over 5-year periods (last bar is a 6-year average). The horizontal black line shows the long-term (entire period) average of 20 days. The number of extremely hot days in Texas was mostly above average between 1910 and 1960 and has been below average since the 1960–1964 period, with the exception of the 2010–2014 period. Sources: CISESS and NOAA NCEI. Data: GHCN-Daily from 24 long-term stations.




Figure 3: Observed annual number of very warm nights (minimum temperature of 75°F or higher) for Texas from 1900 to 2020. Dots show annual values. Bars show averages over 5-year periods (last bar is a 6-year average). The horizontal black line shows the long-term (entire period) average of 20 days. The 1970s saw a record low number of very warm nights. That number increased in the early 21st century, with the record highest number occurring in the 2010–2014 period. Sources: CISESS and NOAA NCEI. Data: GHCN-Daily from 24 long-term stations.




Figure 4: Observed (a) annual number of entire days below freezing (maximum temperature of 32°F or lower), (b) total annual precipitation, (c) annual number of 3-inch extreme precipitation events (days with precipitation of 3 inches or more), and (d) total number of hurricane events (wind speeds reaching hurricane strength somewhere in the state) for Texas from (a, c, d) 1900 to 2020 and (b) 1895 to 2020. In Figures 4a, 4b, and 4c, bars show averages over 5-year periods (last bar is a 6-year average), dots show annual values, and the horizontal black lines show the long-term (entire period) averages: (a) 2.4 days, (b) 27.2 inches, (c) 0.8 days. In Figure 4d, bars show totals over 5-year periods (last bar is a 6-year total). The number of entire days below freezing was above average in the 1970s and 1980s, but since then it has been mostly below the long-term average. Annual precipitation has varied year to year but was well above average during the 2015–2020 period. The number of extreme precipitation events was above average during the 1990s and early 2000s and again in the 2015–2020 period. A typical reporting station experiences about one 3-inch precipitation event per year. There is no long-term trend in the number of hurricanes. Sources: CISESS and NOAA NCEI. Data: (a) GHCN-Daily from 24 long-term stations; (b) nClimDiv; (c) GHCN-Daily from 47 long-term stations.

caused the largest U.S. evacuation in history, Hurricane Harvey is the costliest hurricane in Texas history, with an estimated \$136 billion in damages. Storm surges between 11 and 13 feet along the Texas coast typically have return periods of 25 years (Figure 7). Over the past 30 years (1991–2020), Texas has averaged 149 tornadoes and 4 tornado fatalities per year. Events can occur all year, though activity typically peaks between April and June.

Under both higher and lower emissions pathways, historically unprecedented warming is projected by the end of this century (Figure 1). However, a large range of temperature increases is projected under both pathways, and under the lower pathway, a few projections are only slightly warmer than historical records. Increases in the number of extremely hot days and decreases in the number of

extremely cold days are projected to accompany the overall warming. By 2055, an estimated increase of 20–30 days with temperatures higher than 95°F is projected under one pathway, with the greatest increase in southwestern Texas.

Future changes in annual average precipitation are generally projected to be small (Figure 8), but an increase in extreme precipitation is likely. Furthermore, even if average precipitation does not change, **higher temperatures will increase the rate of soil moisture loss, likely leading to more intense naturally occurring droughts.** Longer dry spells are also projected.

Increased drought severity and increased human demand for surface water will cause changes in streamflow, with extended reductions of freshwater inflow to Texas bays and estuaries. These changes in streamflow will cause temporary or permanent changes to bay salinity and oxygen content, which will have potentially major impacts on bay and estuary ecosystems, such as negatively affecting organism growth, reproduction, and survival.

Future changes in the frequency and severity of tornadoes, hail, and severe thunderstorms are uncertain. However, **hurricane intensity and rainfall are projected to increase for Texas as the climate warms.**

Hurricane Harvey Rainfall, August 24–31, 2017




Figure 5: Rainfall totals in southeast Texas from Hurricane Harvey for August 24–31, 2017. Large areas received more than 30 inches of rainfall, with more than 50 inches in a few locations. The Houston metro area experienced massive flooding, displacing tens of thousands of residents and damaging or destroying more than 100,000 homes and businesses. Source: Martinaitis et al. 2021, Figure 12a. © American Meteorological Society. Used with permission.

Since 1900, global average sea level has risen by about 7–8 inches. It is projected to rise another 1–8 feet, with a likely range of 1–4 feet, by 2100 as a result of both past and future emissions from human activities (Figure 9). Sea level rise has caused an increase in tidal floods associated with nuisance-level impacts. Nuisance floods are events in which water levels exceed the local threshold (set by NOAA’s National Weather Service) for minor impacts. These events can damage infrastructure, cause road closures, and overwhelm storm drains. As sea level has risen along the Texas coastline, the number of tidal flood days has also increased, with the greatest number occurring in 2020 (Figure 10). Future sea level rise will increase the frequency of nuisance flooding and the potential for greater damage from storm surge.




Figure 6: Time series of the Palmer Drought Severity Index for Texas from the year 1000 to 2020. Values for 1895–2020 (red) are based on measured temperature and precipitation. Values prior to 1895 (blue) are estimated from indirect measures such as tree rings. The variances between these data and methodological differences. The fluctuating black line is a running 20-year average. Periods of drought are common in Texas; the most severe droughts since 1895 were those in 1956 and 2011. Prior to 1895, droughts of the 1956 and 2011 severity occurred occasionally. Sources: CISESS and NOAA NCEI. Data: nClimDiv and NADAv2.

Northeast Texas Coastal Surge Return Periods




Figure 7: Storm surge heights occurring at selected average return intervals (10, 25, 50, and 100 years) along the northeast coast of Texas. Sources: CISESS and NOAA NCEI. Data: Needham et al. 2012.

Projected Change in Annual Precipitation




Figure 8: Projected changes in total annual precipitation (%) for the middle of the 21st century compared to the late 20th century under a higher emissions pathway. Hatching represents areas where the majority of climate models indicate a statistically significant change. Texas is part of a large area in the southwestern and central United States with projected decreases in annual precipitation, but most models do not indicate that these changes are statistically significant. Sources: CISESS and NEMAC. Data: CMIP5.

Observed and Projected Annual Number of Tidal Floods for Port Isabel, TX




Figure 10: Number of tidal flood days per year at Port Isabel, TX, for the observed record (1944–2020; orange bars) and projections for 2 NOAA (2017) sea level rise scenarios (2021–2100): Intermediate (dark blue bars) and Intermediate-Low (light blue bars). The NOAA (2017) scenarios are based on local projections of the GMSL scenarios shown in Figure 9. Sea level rise has caused a gradual increase in tidal floods associated with nuisance-level impacts. The greatest number of tidal flood days (all days exceeding the nuisance-level threshold) occurred in 2020 at Port Isabel. Projected increases are large even under the Intermediate-Low scenario. Under the Intermediate scenario, tidal flooding is projected to occur nearly every day of the year by the end of the century. Additional information on tidal flooding observations and scenarios is available at <https://statesummaries.ncics.org/technicaldetails>. Sources: CISESS and NOAA NOS.

Observed and Projected Changes in Global Sea Level




Figure 9: Global mean sea level (GMSL) change from 1800 to 2100. Projections include the six U.S. Interagency Sea Level Rise Task Force GMSL scenarios (Low, navy blue; Intermediate-Low, royal blue; Intermediate, cyan; Intermediate-High, green; High, orange; and Extreme, red curves) relative to historical geological, tide gauge, and satellite altimeter GMSL reconstructions from 1800–2015 (black and magenta lines) and the very likely ranges in 2100 under both lower and higher emissions futures (teal and dark red boxes). Global sea level rise projections range from 1 to 8 feet by 2100, with a likely range of 1 to 4 feet. Source: adapted from Sweet et al. 2017.

Technical details on observations and projections are available online at <https://statesummaries.ncics.org/technicaldetails>.